Enhancing Biological Control in Western Orchards



CONTENTS BACKGROUND 2

Objective I. PESTICIDE EFFECTS ON BC AGENTS...... 3

Studies to estimate the effects of pesticides on BC agents

Objective 2.

Objective 3. SAMPLING TOOLS FOR

Objective 4. WHICH BC AGENTS ARE IMPORTANT?12

Using video and molecular techniques to pinpoint natural enemies of codling moth

Objective 5.

ECONOMIC VALUE OF BC

Evaluating the true value and cost of using enhanced BC

Objective 6.

INFORMATION TRANSFER......15

Providing information to stakeholders in the best way

Year One Progress Report

Project Goals

We have two major goals for this project. The first is to improve the long-term sustainability of the apple, pear and walnut industries in the western US by enhancing biological control of pest insects and mites. The second goal is to synthesize the information developed in this project along with existing information to provide the outreach tools needed to bring about change in grower practices.

Specific Objectives

- 1. Evaluate the sublethal effects of newer pesticides on key natural enemies in laboratory and field assays in apple, pear, and walnut orchards.
- 2. Characterize the phenology of key natural enemies, including timing of emergence from overwintering areas, entry into orchard, and development within the orchard.
- 3. Evaluate attractants as natural enemy monitoring tools and compare them to traditional methods.
- 4. Develop molecular and video methods to monitor predation of codling moth by generalist natural enemies.
- Conduct economic analyses to determine long-term costs associated with IPM programs with and without various levels of biological control.
- 6. Survey clientele to identify optimal ways to present information that will lead to quicker adoption of new technologies; synthesize existing and new information to provide real-time support for pest control decisions by stakeholders.

Project Directors

Washington State University Vincent P. Jones (Project Director) Jay F. Brunner (Executive Committee) Elizabeth H. Beers Karina Gallardo Jessica Goldberger

USDA-ARS Wapato David R. Horton Thomas R. Unruh

University of California, Berkeley Nicholas Mills (Executive Committee)

Oregon State University Peter W. Shearer Steve Castagnoli

Advisory Group

Our advisory group has three outside scientists, one extension agent from each state, and two industry stakeholders from each state.

California

Mike Devencenzi Carolyn Pickel (UC IPM Extension) Jed Walton Marshall Johnson (outside scientist)

Oregon

Rich Garvin Bruce Decker Phil Van Buskirk (OSU Extension)

Washington Dan Flick Nick Stephens Karen Lewis (WSU Extension) Doug Walsh (outside scientist)

Canada

Gary Judd (outside scientist)

INVESTOR NEWSLETTER ISSUE N°3

CODLING MOTH DRIVES IPM IN MULTIPLE CROPS

If not controlled, codling moth (CM) can devastate apple, pear, or walnut crops in the western US (right top row). Because of its damage potential, CM is considered the key pest around which management is focused. With the advent of mating disruption, the intensity of the chemical program has been reduced, allowing biological control (BC) of our secondary pests. The Federal Food Quality Protection Act of 1996, has also resulted in elimination or restriction of organophosphate materials (particularly Guthion or azinphosmethyl in 2012), and indirectly stimulated the registration of a range of new insecticidal chemistries. While these new chemicals have lower mammalian toxicity profiles, their impacts on natural enemies has not been well studied and the change to the new codling moth controls has resulted in outbreaks of secondary pests that were previously controlled by natural enemies.

CAST OF CHARACTERS PESTS AND THEIR NATURAL ENEMIES

Natural enemies can be broadly classed as parasitoids or predators. Parasitoids are intimately associated with a given stage of the pest, making them relatively easy to sample and collect. However, predators are free-living and may or may not be found in tight association with any specific pest; they often feed on a broad range of prey and they may consume part or all of their prey leaving no sign of their activity. Our project is specifically aimed at quantifying the importance of predators as well as conserving all natural enemies by using least disruptive chemicals applied at times when they will have minimum effect on the natural enemy complex. Codling moth damaged apples (L) and adult codling moth (R



Obliguebanded Leafroller attacked by Colpoclypeus florus (L) and Nemorilla pyste (R)



Green apple aphid colony being attacked by Syrphid larva (L), woolly apple aphid being parasitized by Aphelinus mali (R)





European red mite being attacked by *Galendromus occidentalis* (L) , walnut aphid being parasitized by *Trioxys pallidus* (R)



Deraeocoris brevis feeding on pear psylla nymph (L) , aphid being eaten by a lacewing larva (R) (



MORE REALISTIC

IMPROVING ON AN OLD STANDARD

Traditional bioassays treat a pest or natural enemy with a pesticide topically or via residue, then measure acute mortality at some fixed time limit. Our assays are done in two stages (1) "traditional" and (2) "demographic". The demographic assays incorporate multiple modes of pesticide uptake, then looks at the sub-lethal effects (reduced fecundity, altered sex ratios, etc.) and uses demographic models to measure change in population growth rates. This gives us a standard by which to compare different pesticides across natural enemy species.

This set up is more difficult, and the assay arena has to be tailored for each natural enemy and the natural enemy must survive for longer periods than normally used. Each lab has had to develop its own assay arena for each natural enemy evaluated. The figure to the upper right shows the components of the arena used for *Deraeocoris brevis* in Oregon, the assembled arena is shown below it.



Effects on Natural Enemy Population Growth I. Direct and Indirect Effects of Pesticides

This objective has two sub-objectives: 1.1. Laboratory bioassays to identify distinct lethal and sublethal components on one or more life stages of target natural enemies.

1.2. Large-block replicated field studies to verify effects on secondary pests and on natural enemy populations and communities in commercial orchards.

Year 1 Milestones

Complete laboratory bioassays for 2 pesticides for each of the 8 natural enemies.

Summary Objective 1

We have completed the traditional bioassays for more than two materials at each site. The demographic studies have begun at all sites, with most of the *G. occidentalis* studies being completed (Beers lab). Beers also started on objective 1.2 to provide the other labs with suggestions as to problems and solutions for the field trials.

Progress This Year

1.1. Laboratory Bioassays. The acute and sublethal toxicity of five insecticides

(Cyazypyr, rynaxypyr [Altacor], spinetoram [Delegate], novaluron [Rimon], and lambda cyhalothrin [Warrior II]) and two fungicides (sulfur [Kumulus] and copper hydroxide [Kocide] plus Manzate) were assayed for the two predators Deraeocoris brevis (Hemiptera: Miridae) and Galendromus occidentalis (Acari: Phytoseiidae) and the two parasitoids Aphelinus mali (Hymenoptera, Aphelinidae) and Trioxys pallidus (Hymenoptera, Braconidae). Each pesticide was tested at the full field rate (100%) and at a dilute rate (10%) in comparison with a control treated with water. Acute toxicity over 48 hours was assessed for adult parasitoids and for both larval and adult stage predators. In addition, sublethal bioassays were conducted for the predators to assess the following life history performance characteristics: mortality/ survivorship, longevity, fecundity, egg viability, F1 juvenile survivorship, F1 juvenile development time and F1 sex ratio. The acute and sublethal bioassays were carried out in glass arenas under constant laboratory conditions to provide a worst case evaluation of the potential of each pesticide to disrupt the life cycle performance of the selected natural enemies. In the case of G. occidentalis it is

not possible to assess individual life history performance characteristics and so as an alternative, the population growth rate was monitored in replicated mesocosms at the 10 and 100% field rate for each pesticide.

While the laboratory bioassays are still ongoing, example results provide some initial indications of the disruptive effects of the selected pesticides. While Altacor, Delegate and Rimon had no acute effects on larval G. occidentalis, the two latter materials did prove toxic to adults. Warrior proved to be acutely toxic to both nymphs and adults of D. brevis at both rates tested while Rimon proved to be acutely toxic to nymphs of D. brevis (at both rates). For the two parasitoids the acute mortalities were similar to those found for the predators (see Figs on next page), with Delegate and Warrior being most disruptive, Rimon being intermediate and Altacor and Kumulus having minimal impact. Results from the sublethal bioassays are, in most cases, yet to be completed and analyzed. However, results for G. occidentalis and its twospotted spider mite (TSSM) prev are shown (above on the next page) and illustrate the use of the demographic technique.

I.I. TRADITIONAL ASSAYS

SIMILARITIES AND DIFFERENCES

Aphelinus mali (top right) is the parasitoid of the woolly apple aphid and *Trioxys pallidus* (middle right) is the parasitoid of the walnut aphid. Although exposed to completely different pesticide pressures in apple and walnut, they show similarities in response to several of the materials evaluated. Delegate was highly toxic to both parasitoids and Altacor was relatively non-toxic. Rimon affected both parasitoids at the field rate, and Warrior caused \approx 40% mortality for *A. mali*, but 100% for *T. pallidus*.

As more of the assays are completed, we will be able to use data on natural enemy phenology (objective 2) to help determine which materials might be least disruptive at different times of the season.

DEMOGRAPHIC STUDIES

The mesocosm experiments for *G. occidentalis* are the furthest along (right, bottom). We found that Delegate suppressed the growth rate of both *G. occidentalis* and its twospotted spider mite (TSSM) prey. Altacor did not significantly suppress *G. occidentalis*, while Rimon appeared to reduce TSSM and allow reasonable *G. occidentalis* population growth. Belt did not suppress growth rates of *G. occidentalis* and TSSM. Kumulus reduced growth rates of *G. occidentalis* and allowed TSSM populations to increase. Kocide + manzate reduced growth rate of *G. occidentalis* compared to the untreated control while TSSM populations were unaffected.



To verify the outcome from the laboratory bioassays under field conditions and to test methods for next year when more of the large plot studies will be performed in all areas, a pilot study was conducted in two Washington apple orchards this year. Two sites (20-25 acres each) were chosen, one with a history of woolly apple aphid and mite problems, and one without. Four treatments were applied, comprised of different combinations of insecticides used for the first generation of codling moth: either Intrepid or Rimon followed by either Altacor or Delegate. Each treatment was replicated four times, with individual replicates of 1-1.5 acres in size. The three application timings were petal fall (100 CMDD [codling moth degree days]), 1st cover (350 CMDD), and 2nd cover (18 d after 1st cover). Pest and natural enemy populations were sampled at 2-3 week intervals from May through October. Results are shown on the next page in the top box (Field validation).

Next Year

Our plan is to complete lethal and sublethal bioassays for an additional set of natural enemies including the insect predators *Chrysoperla carnea* (Neuroptera.: Chrysopidae) and *Hippodamia convergens* (Coleoptera.: Coccinellidae), and the spiders *Cheiracanthium mildei* (Miturgidae) and *Misumenops lepidus* (Thomisidae). During the 2010 field season, we will also begin replicated field studies of the influence of pesticides that have shown disruptive effects to our natural enemies in the laboratory bioassays in both pear and walnut orchards in addition to apple orchards. The impacts on selected pests and their natural enemies will be monitored.

Implications for the Industry

These laboratory and field assays will allow us to recommend IPM programs that



OBJECTIVES I & 2

I.2. FIELD VALIDATION

VERIFYING LAB DATA

At the Bridgeport orchard, the plots initially treated with Intrepid had consistently lower woolly apple aphid (WAA) densities than those treated with Rimon. The trends within the Intrepid and Rimon plots were similar in that with the second treatment, Altacor treated plots had lower populations than the Delegate treated plots. Overall, it appears that the first spray is most important in "setting the stage" for BC, although the second spray if too disruptive can overcome much of the advantage of having a soft petal fall spray.

The effects of the different pesticides were also evidenced in the inverse relationship of aphid colonies and number of earwigs caught in the different plots. If the pesticides were not suppressing earwig populations, you would expect that the highest earwig numbers would be found in the plots with the highest numbers of WAA colonies, but the opposite trend was seen.

enhance biological control by minimizing disruption of the natural enemy community in orchards. The results of this research will be added to the WSU-DAS and UC-IPM web sites for easy access and will be very useful to apple, pear and walnut growers and pest control advisors. We expect these recommendations to lead to increased biological control in our orchards, which should reduce pesticide inputs leading to higher grower profits and lower worker safety problems.

Researchers Involved in Objective 1

Washington State University (1.1 and 1.2) Betsy Beers Peter Smytheman Randy Talley Lessando Gontijo (graduate student)

University of California at Berkeley (1.1) Nick Mills Kevi Mace Aviva Goldmann

Oregon State University (1.1) Peter Shearer Kaushalya Amarasekare Amanda Borel USDA-ARS Wapato (1.1) Tom Unruh

Minimizing Treatment Impacts 2. Quantifying BC Agent Phenology

Year 1 Milestones

Evaluate phenology in eight apple and three pear orchards; compare attractant traps to traditional methods; start adult phenology studies in walnuts using attractant traps.

Summary Objective 2

All of the milestones for year 1 have been met. Data analysis is in progress and samples should be completely processed before the start of next season.

Progress This Year

This past year, we collected beating tray samples from eight apple orchards in Washington (four in the Quincy area, four near Yakima), and five (instead of three) pear orchards in Oregon (near Hood River). In all of those orchards plus three walnut orchards in California, we used an attractant discovered in our 2008 studies to capture the green lacewing *Chrysopa*



Pesticide



nigricornis (next page bottom photo). We are still processing the beating tray samples, but in general, we can say that the number and diversity of natural enemies collected is a tiny fraction of what we found in our attractant traps (objective 3).

The lure for *C. nigricornis* performed well at nearly all sites (Table on next page). In all situations, the lure caught lacewings earlier and later, and in dramatically higher numbers than what we captured in beating trays (Table on next page). Although beating trays may have their place for estimating populations of certain natural enemies, it is clear that they can distort the impression of natural enemy importance and phenology. Using beating trays alone, *C. nigricornis* would be considered a rare natural enemy; whereas the truth is that it is very common.

In both Washington and California, we had one orchard each that had a very low trap capture. The Washington orchard was a very small orchard that may have contributed to the low catch. At multiple sites over the past two years in Washington using the lacewing attractant, we have not

2. QUANTIFYING PHENOLOGY

DEPENDS ON HOW YOU SAMPLE

Sampling in tree fruits has long used "beating sheets" (upper right) or "limb tapping" to sample natural enemies and pests. The limbs are struck with a rubber hose to dislodge the insects on to the tray below. While this works for some insects, it has a severe limitation: it is an instantaneous sample that only affects the insects on the limb being sampled at that point in time. Insects that occur high on the tree or that move around in response to temperature or light are poorly sampled. Highly mobile insects such as the green lacewings (right lower) are typically under represented in tray counts. Traps on the other hand, accumulate catch over the entire length of the time they are out, so that insects that have a daily movement pattern are represented in the capture if the lure is attractive. The table below shows an extreme example of how poorly beating sheets perform compared to an attractant trap and how that affects our interpretation of natural enemy phenology.



Comparison of the abundance and phenology of the lacewing *Chrysopa nigricornis* measured in Washington apple orchards and Oregon pear orchards using beat-tray samples and attractant traps during 2009.

	Initial Capture		Last Capture		Total Captured		
Orchard	Beat Tray	Attractant	Beat Tray	Attractant	Beat Tray	Attractant	Crop
1	22-Jul	24-Apr	22-Jul	15-Oct	1	7,158	apple
2	24-Jul	13-May	4-Sep	6-Oct	4	13,770	apple
3	none	18-May	none	6-Oct	0	9,108	apple
4	11-Aug	21-May	8 Sept	19-Oct	5	3,195	apple
5	18-Aug	4-Jun	10 Sept	30-Sep	2	717	apple
Difference	75.4 days		42.6 days		12	33,948	Totals
6	26-May	26-May	26-May	2-Oct	11	1,634	pear
7	13-Aug	1-Jun	13-Aug	24-Sep	4	414	pear
8	15-Jun	26-May	15-Jun	2-Oct	9	1,993	pear
9	4-Jun	1-Jun	4-Jun	16-Sep	3	412	pear
10	28-Sep	1-Jun	28-Sep	10-Sep	1	482	pear
Difference	43 days		73.2 days		28	4,935	Totals

C. NIGRICORNIS

A PROMISING START

A preliminary model from two California walnuts orchards (Mills data) and three Washington apple orchards (Jones' data only) shows that phenology of *C. nigricornis* among sites is very similar on a degreeday basis (right). As additional data from the Horton and Shearer labs are compiled, we will continue the efforts to develop a robust phenology model for this key natural enemy, as well as others.



Implications for the Industry

The lacewing attractant shows how sampling using only beating travs distorts the perception of natural enemy importance and timing. We expect that our attractant studies in conjunction with overwintering band emergence studies, will provide us with a much better understanding of the importance, diversity and phenology of our key natural enemies. When the data are complete, it will allow us to time pesticide applications for the pests so that impacts on natural enemies are minimized. When combined with data on sublethal effects on natural enemies (from objective 1), we should have a much better understanding of ways to enhance BC in our orchard systems.

Researchers Involved in Objective 2

Washington State University Vince Jones Shawn Steffan Callie Baker Tawnee Melton

University of California at Berkeley Nick Mills Kevi Mace Aviva Goldmann

Oregon State University Peter Shearer Kaushalya Amarasekare Amanda Borel Kelly Wallis

seen such low captures. In the three commercial apple orchards near Quincy, this year we captured 30,036 (7,158; 13,770; and 9,108) C. nigricornis. In California and Oregon the trap catches were lower, but still substantial (3,113 in two orchards in California and 4,935 in five Oregon orchards). The differences in trap catch between apple, pear, and walnut suggest that we might need high and low sensitivity lures for different situations. The high trap catch in Washington is clearly excessive for determining population trends and using the lure to evaluate impacts of pesticides on C. nigricornis. Lure longevity and dose of the C. nigricornis attractant is examined further in Objective 3.

The phenology of *C. nigricornis* is still being investigated, but it is clear that it emerges fairly late in the season in Washington apples (mid-May) and about a month earlier in California. Unfortunately, we have not found any studies of the temperature-development of *C. nigricornis* in the laboratory, but a survey of nine different studies of lacewings showed that the lower threshold for development varies from 45-50°F. A preliminary model (above) shows great promise that we will be able to change management programs to preserve this important predator.

In addition to the orchard samples, Horton sampled orchards with bands from four

orchards in the Yakima area and Jones collected from eight sites from Brewster WA to Frenchman Hills. The emergence data has all been collected, but only Horton's has identified all specimens and entered the data into the computer for analysis; Jones' data are still being processed. For this coming year, band data will be available for four apple orchards near Yakima, nine apple orchards from Brewster to Quincy, and five pear orchards near Hood River, OR.

Next Year

The beating tray data will be taken from the same orchards in Washington and Oregon and emergence data from overwintering will be available from 13 orchards in Washington and the 5 pear orchards in Hood River. We will also expand the number of lures being evaluated for season-long phenology of natural enemies based on data collected this year in objective 3. This winter, modeling studies on C. nigricornis will continue with validation being performed using data collected next year. Shearer and Jones have submitted two grants leveraging SCRI funds to begin surveying natural enemies in sweet cherry orchards (three orchards in Washington and three in Oregon) that will be included in model development and validation.

3.1. BUILDING A PRACTICAL LURE

A COMBINATION OF BIOLOGICAL RESPONSE AND PHYSICAL CHEMISTRY

Building a practical lure means that we need to have the release rate nearly constant as temperature changes and a release rate that optimizes trap capture. Each chemical tested has a different release profile (top right) that needs to be assessed and then checked for optimum biological activity (bottom right for *C. nigricornis* lure).

We used polyethylene tubing of different thicknesses and if needed, made the packet out of foil with polyethylene windows of various sizes (below).



USDA-ARS Wapato Dave Horton Tom Unruh Gene Miliczky

Improving Our Understanding of Natural Enemies in the Orchard

3. Evaluating New Monitoring Techniques

This objective has three sub-objectives: 3.1. Evaluate release rates of different attractants to determine longevity and the rate needed for optimal trap catch.

3.2. Evaluate the effectiveness of different types of traps.

3.3. Evaluate traps for monitoring diversity, abundance, and phenology of natural enemies, and demonstrate the utility of



lures for assessing the effects of spray programs on natural enemy abundance in orchards.

Year 1 Milestones

Complete studies on longevity and optimal release rates for eight attractants, evaluate four different trap types for selectivity, determine adult phenology using eight attractants in eight apple orchards, three pear orchards, and three walnut orchards; and test in a subset of these orchards whether codling moth cover sprays affect counts of natural enemies.

Summary Objective 3

All of the milestones for year 1 have been met and exceeded. Data analysis is still ongoing for Oregon and some of the Washington samples taken around Yakima. That data analysis should be completed before the start of next season.

Progress This Year

3.1. Lure Release Rates. We

completed our studies on the longevity and release rates for 14 different attractants this year. We generally were able to evaluate release rate and longevity with four different types of lures (typically heat sealed polyethylene tubing of different thicknesses with a cotton dental wick upon which the attractant is applied -see above) during the spring, summer, and fall. The polyethylene tubing was generally acceptable as a release device (top figure above); our studies showed that release rate for each chemical is a function of tubing thickness and temperature. The temperature effects only appeared to be significant between cooler periods (spring or fall) and the summer; in no cases did we have non-linear release of attractants

within a test (generally 1 month periods) related to temperature. Studies with other dispensers (such as mating disruption dispensers) have already shown that release rate is a function of temperature accumulations and we will be performing that analysis this coming winter.

Although the tubing worked well for the majority of attractants tested, for six compounds we needed to develop various modifications of the dispensers because the lure longevity was too short. In four of those cases, we were able to increase the volume of the attractant that was applied to the wick and obtain the desired longevity. However, for at least two of the compounds, we needed to develop a foil based lure with different size polyethylene windows to regulate the release rates. In all cases, we were able to develop lures that would last >28 days during the summer at levels that were biologically active.

In addition to testing the lure release rates, we also performed field bioassays for a number of different types of lures to evaluate the sensitivity of the insects to release rate. In 2008, we had performed this in one orchard with four rates of seven different lures. In that test, the natural enemy diversity was relatively restricted, but there were no real trends in trap catch that could be related to lure release rates. This past year, we ran three experiments where lure rate was evaluated. The first was to restrict the release rates of our C. nigricornis lure, which catches too many lacewings (see objective 2). In that test, we evaluated a normal 4 mil thick bag, a 6 mil half foil bag, and used gray rubber septa loaded with 1, 10 or 100 μ l (facing page, lower figure blue bars). We found that the reduced release rate did significantly reduce the capture of C. nigricornis, with the 6 mil half foil bag lures catching roughly 40% of the normal lure and the $100 \ \mu$ l septa catching $\approx 25\%$ of the normal lure (facing page lower figure, blue bars). We ran a followup trial because the 100 μ l septa load took roughly 7 days to absorb

into the septa, making it impractical for any normal situation. In the second trial, we evaluated the different size windows, which allowed us to adjust the release rate enough to bring the trap catch down to a reasonable level (Facing page lower figure, black bars). The final release rate trial evaluated trap catch for our four new candidate lures, but to date we have only processed traps from the first sample date.

3.3 Evaluating Lure Attractiveness.

This sub-objective had four different experiments run this year (1) long-term testing at eight apple sites in Washington (four in the Wenatchee area, four in Yakima), five pear sites in Oregon near Hood River, and three walnut orchards in California; (2) an in-depth study of the interactive effects of three particularly attractive compounds (geraniol, methyl salicylate and 2-phenylethanol); (3) the investigation of new materials for next years studies, and (4) measuring insecticide impacts using our attractant lures.

3.2. TRAP EFFECTIVENESS

TRAP TYPE "TUNES" CAPTURES

The effect of trap type was evaluated with a combination of three lures and an untreated check. We evaluated the standard white delta trap, an orange delta trap, a white sticky card, and a yellow sticky card (above right). We chose lures to evaluate that had shown a broad spectrum of activity so that we could evaluate trap type for a number of taxa. The lures used were: (1) geraniol + 2-phenylethanol, (2) 2-phenylethanol + methyl salicylate, and (3) acetophenone. We present data for a trap type using only the most active lures for a given taxa.

The yellow card was consistently one of the most sensitive for all the natural enemies, edging out all the other traps types for all natural enemies (right). The effects were most marked for the parasitic hymenoptera (data not shown because we caught >360 per trap on the yellow cards), but the trends were mostly consistent for the other taxa as well. The honeybees did not respond to orange delta traps, white sticky traps, or yellow sticky traps, allowing us to use any of those trap types to minimize undesirable honeybee captures.



3.3. DIVERSITY

LURES ATTRACT DIVERSE TAXA

The majority of specimens captured in north-central WA apples were green lacewings (~24%) and parasitic Hymenoptera (~23%) (right). While the percentages represented by other taxonomic groups were much lower, there were actually thousands of individuals trapped in all groups except the *Stethorus* beetles.

PHENOLOGY OF NATURAL ENEMIES

COMPARISONS SHOW IMPORTANT DIFFERENCES BETWEEN GROUPS

In WA apples, using the top three most attractive lures, we were able to track the phenology of lacewings, syrphids, parasitoids, and predatory Hemiptera (right). From our trapping data, the phenologies of these groups appear to be quite variable. Among the lacewings, we found that *Chrysoperla plorabunda* exhibited a different seasonal pattern than *C. nigricornis* (Fig. A right). At this site in Quincy, WA, the initial *C. nigricornis* flight peaked in late-May/ early-June and then dropped off markedly by late June. However, *C. plorabunda* counts were clearly rising as *C. nigricornis* started dropping off. *C. plorabunda* then exhibited two more peaks, one in early July and another in late July.

Syrphid phenology suggested there may be two "pulses" in adult activity—the first in mid-June and the second in mid-July (Fig B, right). Syrphids are important predators of common apple aphids, as well as woolly apple aphids, which contributes substantially to the suppression of these secondary pests.

Predatory bugs, such as *Campylomma* and *Deraeocoris*, are important predators of codling moth eggs and other soft-bodied prey. *Campylomma* numbers spiked in late June, followed by a pulse in *Deraeocoris* numbers from mid- to late July (Fig. C, right). Importantly, the spike in *Deraeocoris* populations was almost perfectly aligned with the second CM generation. In order to preserve the benefits provided by *Deraeocoris* (e.g., suppression of codling moth, obliquebanded leafroller (OBLR), and the multitude of secondary pests), management strategies might consider concentrating codling moth and/or OBLR sprays during their early-season generations.



WHICH COMPONENTS ARE IMPORTANT IN A BLEND?

LURES ATTRACT DIVERSE TAXA

Our long-term trapping data in NC WA showed the blend of geraniol, 2-phenylethanol (PE), and methyl salicylate (MS) consistently captured more lacewings (five species), syrphids (12 species), and parasitoids (19 different families) than any other blend. The question was which components were the different taxa responding to?

A factorial experiment allowed us to look at all possible combinations of attractants. Green lacewings responded to any lure with PE, syrphids responded to geraniol containing lures, and parasitoids responded best to lures containing PE and to a lesser extent MS.



Long-Term Testing. This past year, we examined ten lure blends as potential tools for monitoring natural enemy diversity, abundance and phenology. We looked at 1) differences in attractiveness among blends of seven different attractant compounds, 2) arthropod diversity and the attraction of particular taxonomic groups to certain lure types, and 3) phenological trends in the four most abundant insect groups. At four commercially managed apple orchards (WSU Sunrise and 3 orchards in the Quincy area of WA), we deployed a total of 160 traps (40 traps/orchard × 4 orchards) from late May to early October, all of which were checked and serviced weekly.

Similar trapping protocols were employed in the orchards sampled in Oregon and California. In this report, we will be presenting data and analyses from the north-central WA apple orchards (Jones lab) and California walnut orchards (Mills lab); data from the other sites has not yet been completely processed. For simplicity, analyses of trap-catch data are confined to traps using the top three most attractive lure blends. In evaluating the numbers caught, it is important to understand that if a particular group is found only during a portion of the test period, the overall mean is decreased dramatically, even though large numbers may have been caught during certain periods of the season.

The lures attract a broad diversity of natural enemies and in some cases, herbivorous taxa (e.g., moths, western flower thrips). In general, our lures bring in large numbers of lacewings and parasitic Hymenoptera, with other groups also well represented in some of our blends. Our lures can be used to develop a better understanding of the phenology of target natural enemy groups (box on facing page, lower figure). However, they also demonstrate that the reliance in the past on primarily one method of sampling (the beating trays) has grossly underestimated numbers and distorted our understanding of natural enemy phenology. For example, we found that beat tray sampling suggested that parasitoid densities spiked early and then late in the season, while our attractant-trapping indicated that these tiny wasps remain abundant throughout the growing season. Designing management plans without a true picture of the phenology of these organisms cannot be successful.

Evaluation of New Attractants. Large numbers of certain parasitoid families (Scelionidae, Eulophidae, Pteromalidae, Eucoilidae, and Trichogrammatidae) were caught with phenyl-acetaldehyde (PAA) lures. The compound cis-jasmonate (CJ) was attractive to two taxonomic groups: syrphids and *Campylomma* (predatory bugs).

Acetophenone (AP), consistently caught a particular large Ichneumonid wasp. Ichneumonids are known to be important enemies of obliguebanded leafroller (OBLR) and codling moth (CM), suggesting that we may have serendipitously discovered an effective lure for this wasp species. AP was also very attractive to western flower thrips. AP is very promising: in an experiment conducted earlier in the season, a blend of AP, PE, and acetic acid was tested against our "best blend" (GER +PE+MS), and the AP blend caught as many lacewings, syrphids, and parasitic wasps as the GER blend.

Next Year

Promising compounds, such as PAA, AP, and CJ, will be incorporated into the general pool of attractants to be tested in the 2010 growing season. These new

OBJECTIVES 3 & 4

INSECTICIDE EFFECTS ON TRAP CAPTURE

Lures Estimate Impact

Our attractant trapping has made it possible to assess more efficiently and accurately the effects of insecticide applications on natural enemies. In one orchard near Quincy, we found that lacewing populations experienced oscillations but never "crashed" (right, upper). This orchard had relatively few sprays applied, and none were organophosphates. Conversely, a different orchard in the same area saw its lacewing population decline sharply and remain low for approximately four weeks, during which two applications of azinphosmethyl (Guthion) were applied (right, lower). During this same time period at other Quincy orchards, our trapping indicated that lacewing populations tended to rise. These findings provide evidence for the common contention that certain insecticides interfere with natural enemy abundance and/or activity.



compounds will also be integrated into investigations of trap type and dose response effects.

Implications for the Industry

In the absence of an easy monitoring tool only pest activity is currently monitored in western orchards. These data will be used to develop an effective monitoring tool for a range of key natural enemies in western orchards for use by growers and pest control advisers (PCAs). A natural enemy monitoring trap will allow growers and PCAs to track changes in natural enemy activity that might be caused by use of disruptive pesticides or unusual climatic conditions. We expect natural enemy traps to become a valuable tool to verify the activity of natural enemies and to enhance recognition of the importance of biological control in our orchards. In turn, this should reduce pesticide inputs leading to higher grower profits and lower worker safety problems.

Researchers Involved in Objective 3

- Washington State University (3.1, 3.2, 3.3) Vince Jones Shawn Steffan
- Callie Baker Tawnee Melton

University of California at Berkeley (3.3) Nick Mills Kevi Mace Aviva Goldmann

Oregon State University (3.3) Peter Shearer Kaushalya Amarasekare Amanda Borel Kelly Wallis

USDA-ARS Wapato (3.3) Dave Horton Tom Unruh Gene Miliczky

Which Predators are Eating CM? 4. CM Predators

The specific sub-objectives are: 4.1. Use digital-video recorders and lowlight capable cameras to detect predation events and to identify key predator species.

4.2. Use DNA and antibody-based methods to detect codling moth remains in predator guts and describe seasonal predatory activity for key predators.

Year 1 Milestones

Complete video monitoring in 2 apple and 1 pear orchards, and complete development of CM antibody, begin generalized collection of natural enemies for later gut content analysis from apple, pear, and walnut.

Summary Objective 4

The video monitoring was problematic and did not give data that was useful in supplementing the molecular gut content analysis. Pitfall trap collections were made from apple and pear orchards, but not from walnut. The LAMP (loop-

4.1. VIDEO EVIDENCE

NOT AS SIMPLE AS IT LOOKS

The video cameras proved unreliable in the field for both technical and biological reasons. To conserve recorder memory, we used motion detection to start recording, but blowing leaves and light shadows caused by moving leaves in the tree canopy triggered recording reducing efficiency. Bird and rodent predators were responsible for virtually all predation events observed (all but 2 examples of ant predation) (left). We are uncertain if the lack of observed insect predators was a result of the artificial observation arenas and set up or truly reflects normal situations in those orchards. We did attempt to use a cover to reduce bird predation and a screen fence to reduce rodents. The cover eliminated birds, but the screen fence (I meter high) did not prevent rodents (*Peromyscus* spp.) from entering the arena.



mediated DNA amplification) procedure was validated and will be used to screen specimens for presence of codling moth.

Progress This Year

4.1. Videos. We evaluated the usefulness of video recording of sentinel codling moth larvae cocoons in one pear and one apple orchard at the USDA-ARS Moxee Farm. Two video recorders each with 4 cameras were tested for 3 months.

4.2. Molecular Methods. A new DNA amplification method was developed and optimized to detect codling moth DNA in the guts of predators. Predators were collected by pitfall trapping at orchards in Yakima and Wenatchee WA and Hood River OR from July through September.

We developed loop-mediated DNA amplification, LAMP, to detect DNA

remains of codling moth (*Cydia pomonella*) in whole-body homogenates of arthropod predators. Six LAMP primers totaling 157 bp were designed within the 226 bp region of an odorant receptor gene of *C. pomonella.* LAMP specificity was challenged with DNA from 17 non-target arthropods including closely related Lepidoptera, and proved to be specific to codling moth (figure below). We used the LAMP procedure to demonstrate detection of *C. pomonella* DNA up to 96 hr postingestion in the crab spider, *Misumenops lepidus.*

We have begun processing the pitfall trap data and expect to finish this winter.

Next Year

We will continue sampling with pitfall traps and will also use beat trays to collect predators for gut content analysis. Video cameras will be used in the lab to observe ground predator behavior with cocooned versus active larvae to determine if their behavioral repertoire supports attack of the cocooned larval stage. Predators collected in pitfall traps will be analyzed for codling moth in gut contents.

Implications for the Industry

The gut content analysis will be used to help clarify which natural enemies are important for our conservation efforts as well as providing targets for our monitoring efforts.

Researchers Involved in Objective 4

USDA-ARS Wapato Tom Unruh Francisco De La Rosa

4.2. MOLECULAR METHODS

BETTER THAN WE HOPED

PCR based detection of DNA in predator gut contents has been worked out, but requires too much purification before processing the sample. This makes them difficult and costly. We developed a new molecular method (LAMP or loop-mediated DNA amplification) that is much cheaper, quicker, and allows use of whole body homogenates of the predator.





A tiny pest with a big bite

It's nagged U.S. apple growers for decades and has now shut down an entire foreign market

By DAN WHEAT World staff writer

WENATCHEE — It's a moth and it's a worm. It's the worm you think of when you think of "the worm" in an apple, a tree fruit

kanogan counties. what is this little bug that's so power I what does it do? alist says. e codling moth, named after an English ing apple in 1747, is just a tiny worm, less three-fourths of an inch long. But it 67.000-ton ships fro ple think of the worm in the

What is the Cost of Enhanced BC? 5. Economic Analysis

Year 1 Milestones

Develop a model to define cost structures of alternative pest treatments to be conducted in the field trials of Objective 1; gather data to obtain preliminary cost structures. Conduct a survey to elicit farmers' willingness to pay for IPM indirect benefits.

Summary Objective 5

Most of year 1 has been spent developing the basic economic models that will be paired with pest control risk/cost scenarios that will be the basis for determining the costs associated with conventional and enhanced BC programs.

Progress This Year

While results from objective 1 have yet to provide a complete picture of the impacts of different pesticides on natural enemies, the results so far show differences that will help us choose IPM scenarios that would be expected to disrupt or conserve biological control in orchard systems. The development of an economic model for apple is in process and will be available for testing the nine different IPM scenarios in the spring of 2010. The nine scenarios assume different CM densities (risk of crop injury), and a response using control options composed of different combinations of pesticides. These scenarios represent a spectrum of

ng moth can lay up to 100 eggs. Larvae do the real da he apple," that people think of, notes tree fruit specialis An adult female codlin

programs that are expected to have varying degrees of impact on biological control in apple orchards. Economic data for the nine scenarios have been gathered and will serve as inputs for the economic model. The economic model will provide a basis for evaluating the relative impacts of different IPM programs on biological control once data from objective 1 are provided in year 2.

The economic benefit of different IPM scenarios will not come just from enhanced biological control, but can occur from the more efficient use of farm labor due to the safety of certain pesticides that result in shorter re-entry periods. The value of the economic model is that results can be generated over more than one growing season and therefore capture the accumulated benefits of IPM programs that might not show an economic benefit if results from only a single year are examined.

A literature review of past studies has been conducted to formulate a model to include growers' willingness to pay for IPM indirect benefits. Most studies used contingent valuation (CV) to elicit grower's willingness to pay for decreased environmental damage by implementing novel technology. We focused on studies dealing with non-point source water pollution abatements. The fact that

5. COST OF ENHANCED BC

A BROAD PERSPECTIVE **ON COST-BENEFITS**

Typical economic studies on the use of biological control focus solely on the number of pesticides reduced or the value of the commodity saved. However, costs associated with just reduced numbers of sprays ignore safety, and markets opened by eliminating pesticide residues.

adoption of IPM programs relying on biological control is a sequential task rather a one-time task, will make it difficult to assess a one-time value for decreased water contamination, as it usually is the CV approach. Thus, we decided to not use the willingness to pay format to elicit the value for indirect benefits of IPM programs. We will be obtaining these parameters from previous related studies and/or may opt a new assessment tool (Specialty Crops (SISC) Pesticide Metric) that is currently being developed.

A survey is being developed under objective 6 that is coordinated with this objective. The initial survey will be conducted in the walnut industry in CA. This survey will include questions that will help us to understand the likelihood of adopting biological control in orchard systems and will serve as a model for surveys to be conducted in pome fruit in subsequent years of this project. The walnut survey will be conducted in the winter of 2009-10 and results will be available in the spring of 2010.

Next Year

Plans for year 2 are to apply the economic model to apple walnut and pear IPM scenarios using data generated from surveys and objective 1.

OBJECTIVES 5 & 6



Implications for the Industry

The kinds of data generated from economic analysis will inform growers and consultants of the tradeoffs of using different IPM strategies that act to conserve biological control in orchards. The outcomes can be used to design education programs that links to the negative impacts of new insecticides on natural enemies based on objective 1.

Researchers Involved in Objective 5

Washington State University Karina Gallardo Jay Brunner

Optimizing Technology Transfer 6. Outreach

There are three sub-objectives: 6.1. Survey clientele to identify optimal ways for presenting information to allow quicker adoption of new technologies and speed technology transfer.

6.2. Synthesis of existing information and information developed in this project into new IPM programs that will lead to enhanced biological control.

6.3. Develop an educational program for the industry to help adoption of newly developed IPM programs.

Year 1 Milestones

This part of the grant was initially intended to start in year 2 and run through the end of the grant. However, we strongly felt the need to move on this area and have made significant progress in parts of the future year's milestones.

Summary Objective 6

Year 1 was not considered to be the start of this section of the project. However, we have designed surveys that will be used for walnut growers, then adapted to apple and pear growers. We have also started giving presentations, preparing papers, and have set up a project web site that details progress and provides general information to interested stakeholders. Progress reports will also be posted on the site.

Progress This Year

6.1. Survey. A survey of walnut growers in California using the Tailored Design Method is in the final stages of preparation. The project personnel will work with the WSU Social and Economic Sciences Research Center (SESRC) to implement the California walnut grower survey in January-March 2010. During spring/summer 2010, a preliminary analysis of survey results will be conducted and reports prepared. Results from the walnut survey will inform the structure of future surveys of pear growers in Oregon (Winter 2010/11) and apple growers in Washington (Winter 2011/12).

The walnut survey includes questions about pest management practices, sources growers access for information, knowledge

6. OUTREACH

SPREADING THE NEWS

Although we had planned no milestones for this objective in year 1, we have been active discussing preliminary research findings and giving growers information on ways to enhance biological control. We have given multiple presentations at field days, developed web pages, handouts, and articles in grower publications.

of biological control, constraints to the adoption of biological control, research and outreach needs, farm characteristics, and grower demographics. Using the survey data, we will build multivariate/ econometric models to measure the effects of various factors on the likelihood of adopting biological control. The first step in the survey process (*i.e.*, working with colleagues to draft the California walnut grower survey and obtaining the names and addresses of growers to compile a mailing list for the top walnut-producing counties), is in its final stages.

The walnut survey data will help us better understand farmers' pest management decision-making helping to ensure the effectiveness and speed of the transfer of biological control (and related IPM) knowledge and technology to growers. Additionally, survey results will inform future biological control research, education, and outreach activities.

6.2. *New IPM Programs.* The work in this area is targeted for years 3-5 of the project. We will discuss and begin to organize data from the project this winter. This will help with coordination of data collection and reporting so that synthesis and incorporation flows as a natural product of the project.

6.3. Education Program. A web site was established (<u>enhancedbc.tfrec.wsu.edu</u>)

SCRI PROGRESS REPORT

as a primary means of posting information about the project. It has been very useful as a place to refer interested individuals desiring to learn more about the project and will serve as a primary location for posting research results and reports generated by the project. The web site will also eventually serve as a location for educational materials developed by project.

The project's Advisory Committee will be used over the next 1-2 years to provide input in developing a comprehensive educational program for the project. The educational program will be informed from grower survey results.

Implications for the Industry

The primary impact of the project in year 1 has been to raise an understanding of the value of biological control in orchard systems and to explain the projects expected contributions to enhancing biological control in orchards. The survey or walnut growers will provide a framework for future surveys in other crops and help to shape education and outreach efforts in the next four years.

Researchers Involved in Objective 6

Washington State University Jay Brunner Jessica Goldberger Nadine Lehrer Wendy Jones Shawn Steffan Vince Jones

USDA-ARS Wapato Tom Unruh

Project Output

Presentations at meetings

Beers, E. H. 2009. Cover crops: Inviting Natural Enemies into Your Orchard . Int. IPM Symposium, Portland, OR 24-26 March 2009. (symposium).

Beers, E. H. 2009. Disruption of Secondary Pests of Apple in the Northwest by Reduced-Risk Pesticides Int. IPM Symposium, Portland, OR 24-26 March 2009. (symposium). Jones VP, JF Brunner, D Horton, TR Unruh, EH Beers, K Gallardo, J Goldberger, NJ Mills, PW Shearer and S Castagnolli. 2008. Enhance BC in Western US (poster). WA State Hort. Assoc. Ann Meeting, Yakima, WA.

Jones VP. Using pest and natural enemy phenology to enhance biological control in orchards. Int. IPM Symposium, Portland, OR 24-26 March 2009. (symposium)

Jones VP, JF Brunner, GG Grove, B Petit, GV Tangren, WE Jones. Evolution and implementation of decision support systems in perennial crops. Pacific Branch Ent. Soc. of Am., San Diego, CA. 31 April 2009. (symposium)

Mills, N.J. 2008. Enhancing biological control to stabilize western orchard IPM systems. Walnut growers meeting, Stockton, CA, Nov 5, 2008.

Mills, N.J. 2009. Biological control in Western tree fruit: opportunities and limitations. 81st Pest and Disease Management Conference, Portland, OR, Jan 14, 2009

Steffan, SA. Biocontrol in Pacific NW Orchards. PMTP meetings in northern WA: May 28, June 3, June 10, 2009.

Steffan, SA. Biocontrol Innovation at the WSU Tree Fruit Research and Extension Center. WSU Field Day, 22 July 2009.

Unruh, T.R. 2009. Enhancing western orchard biological control. WSU Field Day, 15 July 2009

Publications

Jones VP, TR Unruh, DR Horton, NJ Mills, JF Brunner, EH Beers, PW Shearer. 2009. Tree fruit IPM programs in the Western United States: the challenge of enhancing biological control through intensive management. Pest Management Sci. In Focus article. (in press)

Jones VP, JF Brunner, GG Grove, B Petit, GV Tangren, WE Jones. 2010. A webbased decision support system to enhance IPM programs in Washington tree fruit. Pest Management Sci. (in press)

Web pages

enhancedbc.tfrec.wsu.edu (entry to portal)

Extension Handouts

Jones VP and S Steffan. Common natural enemies in Washington tree fruits.

New Proposals Submitted to Leverage SCRI Funding

Jones VP, SA Steffan, JF Brunner. Evaluating and enhancing biological control in organic orchards. Washington Tree Fruit Research Commission. (2 years, \$136,983).

Shearer PW, K. Amarasekare, VP Jones, SA Steffan. Improving biological control of insect pests of cherry. Washington Tree Fruit Research Commission. (2 years, \$79,485).

Jones VP, SA Steffan. Enhancing biological control in cherries. Washington State Commission on Pesticide Registration. (1 year, \$23,189).

Unruh TR, DR Horton, EH Beers. Efficacy and best practicides for predator releases: lacewing, beetles, and mites. Washington Tree Fruit Research Commission (3 years, \$237,997)

We gratefully acknowledge: Matching Fund Sources:

Washington Tree Fruit Research Commission California Walnut Board Washington State University Oregon State University University of California, Berkeley **FIELD PLOTS**

California Growers in Suisun Valley, Davis Oregon Growers in Hood River Washington Growers in Quincy, Bridgeport, Frenchman Hills, Wapato, and Yakima